

The Complete Set of Infinite Volume Ground States for Kitaev's Abelian Quantum Double Models

Matthew Cha

Department of Mathematics, UC Davis

joint work with Pieter Naaijken and Bruno Nachtergaele

QMath13, October 10, 2016

Kitaev's Quantum Double Model (Kitaev, 2003)

Let G be a finite group and \mathcal{B} the bond set of \mathbb{Z}^2 . For $e \in \mathcal{B}$, assign a $|G|$ -dimensional Hilbert space with an orthonormal basis labeled g .

Interaction terms are defined for each **star** v and **plaquette** f by

$$A_v = \frac{1}{|G|} \sum_{g \in G} A_v^g, \quad \text{and} \quad B_f = B_f^e,$$

where

$$A_v^g \begin{array}{c} \uparrow g_4 \\ \xrightarrow{g_1} \quad \xrightarrow{g_3} \\ \uparrow g_2 \end{array} = \begin{array}{c} \uparrow gg_4 \\ \xrightarrow{g_1\bar{g}} \quad \xrightarrow{gg_3} \\ \uparrow g_2\bar{g} \end{array} \quad B_s^h \begin{array}{c} \xrightarrow{g_2} \\ \uparrow g_1 \quad \uparrow g_3 \\ \xrightarrow{g_4} \end{array} = \delta_{h, g_1g_2g_3g_4} \begin{array}{c} \xrightarrow{g_2} \\ \uparrow g_1 \quad \uparrow g_3 \\ \xrightarrow{g_4} \end{array}$$

The local **Hamiltonian** for $\Lambda \subset \mathcal{B}$ a finite subset is:

$$H_\Lambda = \sum_{v \in \mathcal{V}_\Lambda} (I - A_v) + \sum_{f \in \mathcal{F}_\Lambda} (I - B_f)$$

When $G = \mathbb{Z}_2$ this is the well known **toric code** model. Some general properties of the q.d. models include:

- ▶ The interactions terms form a commuting family of projectors.
- ▶ The space of ground states is **frustration-free** and **topologically ordered**, when defined on a surface of genus g the ground state degeneracy only depends on g , local observables cannot distinguish between ground states (local topological quantum order).
- ▶ Excitations occur when one of the f.f. conditions are violated. Ribbon operators generate excitations at their endpoints from a ground state. The excitations are **anyons**, that is, they obey braided statistics. If G is abelian, they can be labeled by a pair $(\chi, c) \in \hat{G} \times G$, where \hat{G} is the group of characters.

Thermodynamic Limit

The quasi-local algebra of **observables** is

$$\mathcal{A} = \overline{\bigcup_{\Lambda \subset_f \mathcal{B}} \mathcal{A}_\Lambda}^{\|\cdot\|} \quad \text{where} \quad \mathcal{A}_\Lambda = \bigotimes_{e \in \Lambda} M_2(\mathbb{C}).$$

The **dynamics** is given by a one-par. group of automorphisms of \mathcal{A} ,

$$\tau_t = e^{it\delta} \quad \text{where} \quad \delta(\cdot) = \lim_{L \rightarrow \infty} [H_{\Lambda_L}, \cdot]$$

where the limit is in the strong sense and Λ_L is any monotone sequence absorbing \mathcal{B} , e.g., Λ_L is the bond set of $[-L, L]^2$. The local algebra of observables, $\mathcal{A}_{loc} = \bigcup_{\Lambda \subset_f \mathcal{B}} \mathcal{A}_\Lambda$ is a core for δ .

A **state** is a linear functional $\omega : \mathcal{A} \rightarrow \mathbb{C}$ such that $\omega(I) = 1$ and $\omega(A) \geq 0$ if $A \geq 0$.

Definition

A state ω is called an infinite volume **ground state** if

$$\omega(A^* \delta(A)) \geq 0 \quad \text{for all } A \in \mathcal{A}_{loc}.$$

Let K denote the set of all infinite volume ground states.

- ▶ This definition expresses that local perturbations do not decrease the energy of a ground state.
- ▶ Infinite volume ground states are often obtained as weak* limit of finite volume ground states. The choice of finite volume **boundary conditions** play a crucial role.

A ground state satisfying the conditions $\omega(A_v) = 1$ and $\omega(B_f) = 1$ for all stars v and plaquettes f is called **frustration-free**.

Theorem (Alicki-Fannes-Horodecki (2007), Naaijken (2011), Fiedler-Naijken (2015))

There exists a unique translation invariant ground state ω^0 of the quantum double model. ω^0 satisfies the following properties:

- ▶ ω^0 is the unique frustration free ground state.
- ▶ ω^0 is a pure state.
- ▶ Let $(\pi_0, \Omega_0, \mathcal{H}_0)$ be a GNS-representation for ω^0 and H_0 be the Hamiltonian in this GNS representation. Then,
 $\text{spec}(H_0) = \mathbb{Z}^{\geq 0}$ with a simple ground state eigenvector Ω_0 .

In particular, ω^0 is a **gapped** ground state.

Superselection Sectors

Now let G be a finite **abelian** group.

Naaijens (2011), Fiedler-Naaijens (2015) constructed **single excitation** states in the thermodynamic limit. These states are labelled by their charge type and position of the charge, $\omega_s^{\chi,c}$.

Furthermore, distinct charge labels correspond to inequivalent states, and hence different **superselection sectors**. The superselection structure is completely described by the representation theory of the q.d., $\text{Rep}(\mathcal{D}(G))$.

The single excitation states $\omega_s^{\chi,c}$ are infinite volume ground states. Indeed, the set of ground states decomposes into $|G|^2$ sectors corresponding to the different charge types.

Theorem (C-Naaijkens-Nachtergaele, 2016)


Let $\omega \in K$ be a ground state of the q.d. model for abelian group G . Then there exists disjoint subsets $K^{\chi,c}$ of K such that ω has the convex decomposition

$$\omega = \sum_{\chi \in \widehat{G}, c \in G} \lambda_{\chi,c}(\omega) \omega^{\chi,c} \quad \text{where} \quad \omega^{\chi,c} \in K^{\chi,c}.$$

Furthermore,

- ▶ For the single excitation states, $\omega_s^{\chi,c} \in K^{\chi,c}$
- ▶ $K^{\chi,c}$ is a face in the set of all states. In particular, if $\omega^{\chi,c}$ is an extremal point of $K^{\chi,c}$ then $\omega^{\chi,c}$ is a pure state.
- ▶ If $\omega^{\chi,c} \in K^{\chi,c}$ is a pure state then it is equivalent to the single excitation state $\omega_s^{\chi,c}$.

The strategy of the proof is to reduce the infinite volume calculation to a finite volume calculation. In particular, we find a boundary term for every box such that the restriction of any infinite volume ground state to the box is a ground state of the finite volume Hamiltonian plus the boundary term.



Concluding Remarks

- ▶ The complete ground state problem has been solved for the XY -chain by Araki-Matsui (1985), for the XXZ -chain by Matsui (1996) and Koma-Nachtergaele (1998), and for the finite-range spin chains with a unique f.f. MPS ground state by Ogata (2016). Our result is the first solution to the ground state problem for a quantum model in two-dimensions.
- ▶ A current challenge in mathematical physics is the classification of gapped ground state phases. One approach is to construct a complete set of invariants. The invariance of the structure of anyon quasi-particles is usually taken as fact. Our results are a first step in rigorously studying the stability properties of the superselection structure of quantum double models.